

英文論文 – 2016年

No	論文名	著者名	雑誌名	発行年及び掲載頁	その他
1	Post-traumatic Stress and Growth Among Medical Student Volunteers After the March 2011 Disaster in Fukushima, Japan: Implications for Student Involvement with Future Disasters.	Anderson D, Prioleau P, Taku K, Naruse Y, Sekine H, Maeda M, Yabe H , Katz C, Yanagisawa R.	Psychiatr Q.	2016 Jun;87(2):241-51.	10.1007/s11126-015-9381-3.
2	Three-year trend survey of psychological distress, post-traumatic stress, and problem drinking among residents in the evacuation zone after the Fukushima Daiichi Nuclear Power Plant accident [The Fukushima Health Management Survey].	Oe M, Fujii S, Maeda M, Nagai M, Harigane M, Miura I, Yabe H , Ohira T, Takahashi H, Suzuki Y, Yasumura S, Abe M.	Psychiatry Clin Neurosci.	2016 Jun;70(6):245-52.	10.1111/pcn.12387.
3	Reduced frontal activation during verbal fluency task in chronic hepatitis C patients with interferon-based therapy as measured by near-infrared spectroscopy.	Abe K, Wada A, Oshima S, Kono S, Takahashi A, Kanno Y, Imaizumi H, Hayashi M, Okai K, Niwa SI, Yabe H , Ohira H.	Hepatol Res.	2016 Apr 7.	10.1111/hepr.12721. [Epub ahead of print]
4	Variants in the DRD2 locus and antipsychotic-related prolactin levels: A meta-analysis.	Miura I, Zhang JP, Hagi K, Lencz T, Kane JM, Yabe H , Malhotra AK, Correll CU.	Psychoneuroendocrinology.	2016 Oct;72:1-10.	10.1016/j.psyneuen.2016.06.002.
5	Association between psychological distress and dietary intake among evacuees after the Great East Japan Earthquake in a cross-sectional study: the Fukushima Health Management Survey.	Uemura M, Ohira T, Yasumura S, Otsuru A, Maeda M, Harigane M, Horikoshi N, Suzuki Y, Yabe H , Takahashi H, Nagai M, Nakano H, Zhang W, Hirosaki M, Abe M; Fukushima Health Management Survey Group..	BMJ Open.	2016 Jul 5;6(7):e011534.	10.1136/bmjopen-2016-011534.
6	Severe Psychological Distress of Evacuees in Evacuation Zone Caused by the Fukushima Daiichi Nuclear Power Plant Accident: The Fukushima Health Management Survey.	Kunii Y, Suzuki Y, Shiga T, Yabe H , Yasumura S, Maeda M, Niwa S, Otsuru A, Mashiko H, Abe M; Mental Health Group of the Fukushima Health Management Survey..	PLoS One.	2016 Jul 8;11(7):e0158821.	10.1371/journal.
7	Decreased calcineurin immunoreactivity in the postmortem brain of a patient with schizophrenia who had been prescribed the calcineurin inhibitor, tacrolimus, for leukemia.	Wada A, Kunii Y, Matsumoto J, Hino M, Nagaoka A, Niwa S, Yabe H .	Neuropsychiatr Dis Treat.	2016 Jul 6;12:1645-50.	10.2147/NDT.S106371.
8	EXTERNAL AND INTERNAL EXPOSURE TO FUKUSHIMA RESIDENTS.	Kamiya K, Ishikawa T, Yasumura S, Sakai A, Ohira T, Takahashi H, Ohtsuru A, Suzuki S, Hosoya M, Maeda M, Yabe H , Fujimori K, Yamashita S, Ohto H, Abe M.	Radiat Prot Dosimetry.	2016 Sep;171(1):7-13.	10.1093/rpd/ncw185.
9	Decreased VEGFR2 expression and increased phosphorylated Akt1 in the prefrontal cortex of individuals with schizophrenia.	Hino M, Kunii Y, Matsumoto J, Wada A, Nagaoka A, Niwa S, Takahashi H, Kakita A, Akatsu H, Hashizume Y, Yamamoto S, Yabe H .	J Psychiatr Res.	2016 Nov;82:100-8.	10.1016/j.jpsychires.2016.07.018.

英文論文 - 2016年

No	論文名	著者名	雑誌名	発行年及び掲載頁	その他
10	Diagnostic accuracy of Japanese posttraumatic stress measures after a complex disaster: The Fukushima Health Management Survey.	脳科学研究戦略推進プログラム/発達障害・統合失調症等の克服に関する研究	Asia Pac Psychiatry.	2016 Aug 9.	10.1111/appy.12248. [Epub ahead of print]
11	Psychological distress of residents in Kawauchi village, Fukushima Prefecture after the accident at Fukushima Daiichi Nuclear Power Station: the Fukushima Health Management Survey.	Yoshida K, Shinkawa T, Urata H, Nakashima K, Orita M, Yasui K, Kumagai A, Ohtsuru A, Yabe H , Maeda M, Hayashida N, Kudo T, Yamashita S, Takamura N.	PeerJ.	2016 Aug 31;4:e2353.	10.7717/peerj.2353.
12	Predictors of severe psychological distress trajectory after nuclear disaster: evidence from the Fukushima Health Management Survey.	Oe M, Maeda M, Nagai M, Yasumura S, Yabe H , Suzuki Y, Harigane M, Ohira T, Abe M.	BMJ Open.	2016 Oct 19;6(10):e013400.	10.1136/bmjopen-2016-013400.
13	The effect of a behavioral activation program on improving mental and physical health complaints associated with radiation stress among mothers in Fukushima: a randomized controlled trial.	Imamura K, Sekiya Y, Asai Y, Umeda M, Horikoshi N, Yasumura S, Yabe H , Akiyama T, Kawakami N.	BMC Public Health.	2016 Nov 8;16(1):1144.	
14	Psychiatric outpatients after the 3.11 complex disaster in Fukushima, Japan	Hori A, Hoshino H, Miura I, Hisamura M, Wada A, Itagaki S, Kunii Y, Matsumoto J, Mashiko H, Katz C, Yabe H, Niwa S,	Annals of Grobal Health	2016 Sep-Oct;82(5):798-805	
15	Prominent increased calcineurin immunoreactivity in the superior temporal gyrus in schizophrenia: A postmortem study.	Wada A, Kunii Y, Matsumoto J, Hino M, Yang Q, Niwa SI, Yabe H .	Psychiatry Res.	2017 Jan;247:79-83.	10.1016/j.psychres.2016.11.018.
16	Mental Health Status of Children After the Great East Japan Earthquake and Fukushima Daiichi Nuclear Power Plant Accident.	Mashiko H, Yabe H , Maeda M, Itagaki S, Kunii Y, Shiga T, Miura I, Suzuki Y, Yasumura S, Iwasa H, Niwa SI, Ohtsuru A, Abe M.	Asia Pacific Journal of Pubric Health	2017 Mar;29(2_suppl):131S-138S.	doi: 10.1177/1010539516675702.
17	Downregulation of GNA13-ERK network in prefrontal cortex of schizophrenia brain identified by combined focused and targeted quantitative proteomics.	Hirayama-Kurogi M, Takizawa Y, Kunii Y, Matsumoto J, Wada A, Hino M, Akatsu H, Hashizume Y, Yamamoto S, Kondo T, Ito S, Tachikawa M, Niwa SI, Yabe H , Terasaki T, Setou M, Ohtsuki S	J Proteomics.	2017 Feb 16. S1874-3919(17)30058-1.	10.1016/j.jprot.2017.02.009. [Epub ahead of print]
18	Evacuation after the Great East Japan Earthquake was associated with poor dietary intake: The Fukushima Health Management Survey.	Zhang W, Ohira T, Abe M, Kamiya K, Yamashita S, Yasumura S, Ohtsuru A, Masaharu M, Harigane M, Horikoshi N, Suzuki Y, Yabe H , Yuuki M, Nagai M, Takahashi H, Nakano H; Fukushima Health Management Survey Group..	J Epidemiol.	2017 Jan;27(1):14-23.	10.1016/j.je.2016.08.002.

英文論文 - 2016年

No	論文名	著者名	雑誌名	発行年及び掲載頁	その他
19	Drinking Behavior and Mental Illness Among Evacuees in Fukushima Following the Great East Japan Earthquake: The Fukushima Health Management Survey.	Ueda Y, Yabe H , Maeda M, Ohira T, Fujii S, Niwa S, Ohtsuru A, Mashiko H, Harigane M, Yasumura S; Fukushima Health Management Survey Group..	Alcohol Clin Exp Res.	2016 Mar;40(3):623-30.	10.1111/acer.12984.
20	Alteration of Duration Mismatch Negativity Induced by Transcranial Magnetic Stimulation Over the Left Parietal Lobe.	Oshima H, Shiga T, Niwa SI, Enomoto H, Ugawa Y, Yabe H .	Clin EEG Neurosci.	2017 Jan;48(1):11-19.	
21	Association between evacuation condition and habitual physical activity in Great East Japan Earthquake evacuees: The Fukushima Health Management Survey.	Nagai M, Ohira T, Yasumura S, Takahashi H, Yuki M, Nakano H, Wen Z, Yabe H , Ohtsuru A, Maeda M, Takase K; Fukushima health Management Survey Group..	Nihon Kosshu Eisei Zasshi.	2016;63(1):3-10.	10.11236/jph.63.1_3. Japanese
22	Decreased 16:0/20:4-phosphatidylinositol level in the post-mortem prefrontal cortex of elderly patients with schizophrenia.	Matsumoto J, Nakanishi H, Kunii Y, Sugiura Y, Yuki D, Wada A, Hino M, Niwa SI, Kondo T, Waki M, Hayasaka T, Masaki N, Akatsu H, Hashizume Y, Yamamoto S, Sato S, Sasaki T, Setou M, Yabe H .	Sci Rep	2017 Mar 23;7:45050.	doi: 10.1038/srep45050
23	Mental Health and Related Factors of Hospital Nurses.	Nukui H, Murakami M, Midorikawa S, Suenaga M, Rokkaku Y, Yabe H , Ohtsuru A.	Asia Pac J Public Health.	2017 Mar;29(2_suppl):161S-170S.	doi: 10.1177/1010539516682589.
24	Changes of Posttraumatic Stress Responses in Evacuated Residents and Their Related Factors.	Oe M, Takahashi H, Maeda M, Harigane M, Fujii S, Miura I, Nagai M, Yabe H , Ohira T, Suzuki Y, Yasumura S, Abe M.	Asia Pac J Public Health.	2017 Mar;29(2_suppl):182S-192S	doi: 10.1177/1010539516680733
25	Exercise Habits Are Important for the Mental Health of Children in Fukushima After the Fukushima Daiichi Disaster.	Itagaki S, Harigane M, Maeda M, Yasumura S, Suzuki Y, Mashiko H, Nagai M, Ohira T, Yabe H ; Mental Health Group of the Fukushima Health Management Survey.	Asia Pac J Public Health.	2017 Mar;29(2_suppl):171S-181S	doi: 10.1177/1010539516686163.
26	The effect of telephone support to evacuees with risks of hypertension and diabetes mellitus after a disaster: the Fukushima Health Management Survey.	Horikoshi N, Ohira T, Yasumura S, Yabe H , Maeda M; Fukushima health Management Survey Group.	Nihon Kosshu Eisei Zasshi.	2017;64(2):70-77.	doi: 10.11236/jph.64.2_70. Japanese.
27	Psychometric Evaluation of the Japanese Version of the Posttraumatic Stress Disorder Checklist in Community Dwellers Following the Fukushima Daiichi Nuclear Power Plant Incident: The Fukushima Health Management Survey	Iwasa H, Suzuki Y, Shiga T, Maeda M, Yabe H, Yasumura S, Niwa O, Matsui S, Ohira T, Mashiko H, Kunii Y, Itagaki S, Miura I, Nakayama Y, Ohta M, Goto A, Hisata M, Kawakami N, Hosoya M, Harigane M, Yagi A, Oikawa Y, Ueda Y, Horikoshi N.	The Fukushima Health Management Survey. SAGE Open.	20160420; 6(2)	doi: 10.1177/2158244016652444.